

Interdisciplinary Treatment for Upper Lip Support with Fixed Implant Supported Prosthesis Using a Sub Nasal Lip Lift

Abstract

Increasing success with endosseous implants in edentulous patients has heightened the interest in using fixed prostheses in this patient population. Despite this trend, upper lip support has been a long-standing challenge, especially in patients with a thin elongated upper lip, termed “senile”, where the presence or absence of a flange compromises the esthetic and functional outcomes. Although, the conventional solution to reestablish elevation and support of the upper lip has been an anterior flange of a removable prosthesis, fixed implant supported prostheses do not incorporate a flange for reasons of hygiene and maintenance. In this case report a patient with atrophic maxillary edentulism and upper lip senility that sought fixed rehabilitation was identified. Following pre-surgical assessment of vertical facial proportions, lip length and morphology, and inter-arch space, a surgical upper lip length reduction was performed using the Subnasal Lip Lift (SNLL) technique. 6 months after the SNLL, when the lip length stabilized, the patient received fixed implant supported monolithic zirconia prosthesis without an anterior flange. The desired amount of lip elevation and tooth display in the final prosthesis was achieved. The lip length remained stable at the 1-year follow-up, Subnasal scars were not evident by 2 months, abnormal lip dynamics were not observed, and improved vermilion eversion was present.

Keywords: Lip support; Fixed implant supported prostheses; Lip lift; Esthetic facial surgery

Case Report

Volume 5 Issue 1 - 2016

Aikaterini Kostagianni^{*}, Todd C Hanna², Daniel A Givan³ and Patrick J Louis⁴

¹Department of Prosthodontics, Tufts University School of Dental Medicine, USA

²Department of Head & Neck Surgical Oncology, Microvascular & Reconstructive Surgery, Oral & Maxillofacial Surgery, USA

³Department of Restorative Sciences, University of Alabama at Birmingham, USA

⁴Department of Oral & Maxillofacial Surgery, University of Alabama at Birmingham, USA

***Corresponding author:** Aikaterini Kostagianni, Department of Prosthodontics, Tufts University School of Dental Medicine, Kneeland Street, Boston MA 02111, USA, Tel: 617-636-0472; Email: katerinakos86@gmail.com

Received: August 05, 2016 | **Published:** August 25, 2016

Introduction

Patients with acquired maxillary edentulism and atrophy often present with facial soft tissue changes that limit the restorative dentist's ability to fabricate a functional and esthetic prosthesis. Often with age, the upper lip becomes thin and elongated, due to atrophy of muscles, fat, and connective tissue [1] and is termed “senile”. Interestingly, the age-related changes of lip do not result in a loss of lip volume, rather the volume is redistributed to increased length resulting in vermilion inversion and ptosis of the lip [2]. One of the negative consequences of the senile lip regards an unpredictable display of oral spaces, such as the buccal corridor and smile line [3], with the maxillary incisal display at rest being decreased or eliminated [4]. These changes, compounded with atrophy of the underlying maxillary skeleton and tooth loss, result in significant labio-facial changes [5]. The conventional prosthetic approach to provide lip support to this patient population is with a denture flange to reestablish lip support and elevation [6]. However, denture flange support occurs by extension of the prosthesis into the vestibule and pushing the lip outward [7,8] which alters muscle pull and subsequent tissue movement. This artificial support pushes the orbicularis oris outward uniformly and allows the superior anterior border of the flange to develop an unaesthetic Subnasal convexity. The lifting effect of the flange also seems inadequate in correcting an inverted vermilion border and does not restore convexity to the philtrum columns. Although

flanges provide support to a senile lip, they cannot be employed with fixed restorations for reasons of maintenance and hygiene. Additionally, the amount and thickness of material required may limit biomechanical function [9,10]. However, many patients desire fixed implant retained prosthetic solutions but lack adequate soft tissue support without a flange. Calvani evaluated lip support of 412 patients with differing prostheses and thin lips and determined 66% of full-arch implant supported fixed restorations presented with a horizontal groove in the upper lip [11]. This groove, termed a “split philtrum line”, can develop with a fixed implant supported prosthetics when there is inadequate support inferior to the nasal base.

Nonetheless, surgical augmentation of the senile upper lip can facilitate the restoring dentist's ability to provide an esthetic, fixed maxillary prosthesis for the edentulous maxilla minimizing the unwanted effects of atrophy, tissue thinning, and loss of support. Known as a Subnasal Lip Lift (SNLL), it effectively shortens the lip while simultaneously increasing the vermilion display by eversion (Figure 1). While the literature supports that the SNLL procedure has predictable esthetic outcomes to improve age-related changes to the upper lip with intact muscular function, the technique has also been refined in the past decades to diminish scar formation, with meticulous closure, laser resurfacing, or dermabrasion [5,9,12-16]. This technique was employed in this case report to demonstrate the application of surgical management of maxillary

edentulism accompanying a senile upper lip that was restored with a flangeless fixed implant-supported maxillary prosthesis. The purpose was to elevate the upper lip to an ideal position

and philtrum contours for esthetic tooth display and lip support, thereby reducing the need for an anterior flange.




Figure 1: Subnasal lip lift procedure to excise tissue at the Nasal base which is then sutured to shorten the lip and avert the vermilion to increase display.

Clinical Report

A 78 year-old female presented to the University of Alabama at Birmingham School of Dentistry seeking care. She expressed dissatisfaction with the function and esthetics of her existing maxillary prosthesis and sought rehabilitation with fixed implant supported restorations. Clinical evaluation revealed maxillary edentulism with moderate atrophy, a senile upper lip with lack of vermilion display and an unnatural convexity caused by her existing removable prosthesis due to the attempt to create an ideal maxillary tooth display (Figure 2). Vertical facial proportions, lip length/morphology and vestibular depth were assessed. Lip length was measured in repose from the Subnasal to the stomion superius and was found 25mm (Figure 3). These findings indicated that patient's desire for maxillary fixed implant supported prosthesis could not be achieved only with prosthetic treatment. Thus, an interdisciplinary treatment plan was developed to reestablish a natural lip support with a combination of prosthetic tooth support and a SNLL procedure. Prosthodontic evaluation included a trial diagnostic tooth set-up without a flange to judge lip support provided by the prosthetic teeth alone and measure the available restorative space [17,18]. The lip contours were visualized without the flange to determine a realistic pre-surgical assessment of the need for additional lip support [19,20]. The teeth were set up to ideal dimensions for a fixed implant supported prosthesis, with approximately 10mm interarch space [18,21]. This set-up was also utilized as a reference to evaluate the desired amount of lip reduction. In final planning with the surgeon and prosthodontist, an 8 implant-supported screw retained monolithic Zirconia restoration was selected.

The Subnasal Lip Lift procedure (SNLL)

The planned lip excision was marked with the patient in the upright position for accurate lip length assessment and prior to administering local anesthesia to avoid tissue distortion (Figure 4). A bullhorn-type incision in the Subnasal region was designed. The superior incision line approximates the naso-labial junction, and extends around the ala-facial groove. The inferior incision line mirrors the pattern superior excision with a separation of the excisions at the desired level of lip lift. The incision is made with a #15 blade and carried into the subcutaneous plane. The underlying orbicularis oris muscle was not excised because of the potential for irregular lip dynamics, however, some advocate muscle excision proposing it offers greater stability of the lift with less relapse [5,22]. Closure was achieved with meticulous interrupted 4-0 Vicryl subcutaneous sutures (Ethicon, Somerville, NJ) and 5-0 nylon skin sutures taking care to avert the wound edges (Figure 5). The wound was treated with topical antibiotic ointment, applied daily for 1 week, and the nylon sutures were removed in 7 days. Sunblock was recommended for six months to avoid any potential for hyper pigmentation. On the same day of the SNLL 8 dental implants were placed utilizing a surgical guide fabricated from the trial teeth set-up (Figure 6). Delayed engagement of the implants was applied due to inadequate primary stability and a tissue-borne interim denture was inserted one week after SNLL. The anterior flange was reduced as much as possible to accommodate the lip elevation and minimize excessive anterior tooth display. It is important to inform the patient that the ideal dimensions and esthetics of this provisional prosthesis may not be feasible during the healing phase. However, if the initial implant stability permits, the denture prosthesis may be immediately converted to a flangeless screw-retained fixed provisional restoration, which will allow for more optimal esthetics.

After approximately six months of healing following the implant placement and the SNLL, an implant supported fixed provisional restoration was fabricated to reassess tooth position, lip support, and to serve as a guide for the final implant supported prosthesis. The delay is not only for adequate osseointegration, but to allow for the anticipated partial relapse of the SNLL prior to stabilization of the lift length [22]. The patient had a preoperative lip length of 25 mm and underwent an 8 mm Subnasal tissue resection resulting in a 19 mm immediately post-operative. At 6-months lip length was found to be 21 mm lip (Figure 7). These measurements were stable at 1-year follow-up, while the Subnasal scar was inconspicuous by 2 months. For the definitive

prosthesis a screw-retained monolithic Zirconia restoration was fabricated; while the prosthesis replaced teeth and tissue, there was no extension into anterior vestibule. The final outcomes demonstrated an increase in vermilion show and elimination of a Subnasal bulge for the upper lip. Satisfactory lip support was accomplished with the combination of the SNLL and the fixed implant-supported restoration. Philtrum columns and an age-appropriate vermilion display were achieved, (Figure 8). At the one-year follow up patient expressed high satisfaction with the esthetic results of her treatment, as well as improved function of her implant-supported prosthesis.



Figure 2: Senile lip without denture flange support noting excess length, flattening of contours, and vermilion inversion (A). Senile lip with a well-constructed denture flange (B, C and D) which uniformly pushes out the lip with loss of philtrum columns, unnatural fullness, inversion of the vermilion, and a subnasal convexity.



Figure 3: Pre-operative assessment of a senile lip with denture removed.



Figure 4: Bullhorn-type SNLL marking are made with the patient upright, prior to local anesthesia.



Figure 5: A Surgical excision of superficial layers without encroachment into the underlying muscle tissue. B. Post-surgical final closure prior to anticipated partial relapse.




Figure 6: Surgical guide duplicated from the diagnostic teeth set-up and placement of 8 dental Implants.



Figure 7: 6-month postoperative assessment with screw-retained implant supported provisional prosthesis. Note that no flange is incorporated for lip support (A. Lateral high smile view, B. Lateral repose view, C. Provisional prosthesis)



Figure 8: One-year outcome with Zirconia fixed implant supported prostheses (A. Frontal high smile view, B. Frontal repose view, C. Monolithic Zirconia screw retained Prosthesis) noting a natural support contour and the presence of age-appropriate philtrum columns and vermilion display.

Discussion

Inadequate lip support may result from reasons other than maxillary atrophy with a senile lip, and a comprehensive oral and facial evaluation should be performed to determine if soft tissue surgical procedures, such as a SNLL, are appropriate rather than procedures involving hard tissues. To determine the etiology, an assessment should include the facial height, facial profile, lip thickness and lip length [10]. The assessment is done with and without patients existing dentures in place, if available, evaluating both the vertical facial proportions of the upper lip, lower lip, and chin as well as the contribution of the vertical dimension of occlusion. Documentation of upper lip length, depth of the vestibule, and height of the anterior maxillary ridge is important to aid interdisciplinary planning for SNLL treatment. The fullness of the upper lip is assessed by a measure of the nasiolabial angle (normal range 85°-105°) or from measures against normal diagnostic lines, such as Burstone's, Steiner's, or Ricket's lines [23]. Factors such as lip thickness, morphology of the nasal base, and presence of facial hair effects the perception of support and is challenged with the senile lip. The normal upper lip length

at rest is estimated to range from 18-22 mm [23,24]. However, rather than endorsing a set numeric range to guide the need for surgical correction, greater emphasis should be placed on proportional values to account for individual and ethnic variance. The vermilion follows proportions by division into thirds with the upper lip being 1/3 and the lower lip being 2/3 of total vermilion height [22,26,27]. These proportions believed to be relatively consistent in classical measures of attractiveness.

Perenack noted the relative contraindications to an SNLL are a short lip length, less than 18 mm, or a history of forming hypertrophic scars and keloids, otherwise excellent outcomes could be routinely obtained. 5 If the lip length is found to be approximately 18 mm or less, the lip may be well supported by the prosthetic teeth alone since their position and angulation are the major supporting features of the inferior portion of the upper lip. The changes in lip length associated with orthognathic surgery are also well established whereas conventional bone grafting of the atrophic maxilla may provide only minimal success in supporting the senile lip largely due to the extent of grafting necessary and does not address the lip length or morphology [27-29]. Evaluation

of the apical position of the upper lip during maximum smile is also critical. Lip function that provides less gingival display was often deemed more esthetic, and a low lip line is advantageous for the fixed anterior prostheses, since the tissue-prosthesis junction can be visible in high smile patients [30,31].

Summary

The described Subnasal lip lift (SNLL) offered a predictable and esthetic improvement of age-related upper lip changes, and allowed for a functionally successful restoration with fixed implant-support prosthetics. Esthetic and functional outcomes were enhanced by normalizing upper lip length, increasing vermilion show, increasing tooth display, eliminating a Subnasal convexity, and reduction of restorative material. Additional factors of expense, multiple staged procedures, and overall treatment time must be considered. Overall, the Subnasal lip lift provided a highly satisfactory treatment outcome. Preservation of muscular function and avoidance of an underlying flange are key elements of this treatment modality.

References

- Perkins SW, Sandel HD (2007) Anatomic considerations, analysis, and aging process of the perioral region. *Facial Plast Surg Clin North Am* 15(4): 403-407.
- Iblher N, Stark GB, Penna V (2012) The aging perioral region--do we really know what is happening? *J Nutr Health Aging* 16(6): 581-585.
- Moore T, Southard KA, Casco JS, Qian F, Southard TE (2005) Buccal corridors and smile esthetics. *Am J Orthod Dentofacial Orthop* 127(2): 208-213.
- Hulsey CM (1970) An esthetic evaluation of lip-teeth relationships present in the smile. *Am J Orthod* 57(2): 132-144.
- Perenack J (2005) Treatment options to optimize display of anterior dental esthetics in the patient with the aged lip. *J Oral Maxillofac Surg* 63(11): 1634-1641.
- Mericske-Stern RD, Taylor TD, Belser U (2000) Management of the edentulous patient. *J Clin Oral Implants Res* 11(Suppl. 1): 108-125.
- Kamashita Y, Kamada Y, Kawahata N, Nagaoka E (2006) Influence of lip support on the soft-tissue profile of complete denture wearers. *J Oral Rehabil* 33(2): 102-109.
- Ushijima M, Kamashita Y, Nishi Y, Nagaoka E (2013) Changes in lip forms on three-dimensional images with alteration of lip support and/or occlusal vertical dimension in complete denture wearers. *J Pros Res* 57(2): 113-121.
- Taylor T (1991) Fixed implant rehabilitation for the edentulous maxilla. *Int J Oral Maxillofac Implants* 6(3): 329-337.
- Bidra A (2011) Three-dimensional esthetic analysis in treatment planning for implant-supported fixed prosthesis in the edentulous maxilla: review of the esthetics literature. *J Esthet Restor Dent* 23(4): 219-236.
- Calvani L, Michalakis K, Hirayama H (2007) The influence of full-arch implant-retained fixed dental prostheses on upper lip support and lower facial esthetics: preliminary clinical observations. *Eur J Esthet Dent* 2(4): 420-428.
- Santanché P, Bonagiggo C (2004) Lifting the Upper Lip: Personal Technique. *Plast Reconstr Surg* 113(6): 1828-1835.
- Yoskovitch A, Fanous N (2003) Correction of thin lips: a 17-year follow-up of the original technique. *Plast Reconstr Surg* 112: 671-672.
- Ramírez OM, Khan AS, Robertson KM (2003) The upper lip lift using the bulls horn approach. *J Drugs Dermatol* 2(3): 303-306.
- Waldman SR (2007) The subnasal lift. *Facial Plast Surg Clin North Am* 15(4): 513-516.
- Penna V, Iblher N, Bannasch H, Stark GB (2010) Proving the effectiveness of the lip lift for treatment of the aging lip: a morphometric evaluation. *Plast Reconstr Surg* 126(2): 83e-84e.
- Ellis E, McFadden D (2007) The value of a diagnostic setup for full fixed maxillary implant prosthetics. *J Oral Maxillofac Surg* 65(9): 1764-1771.
- Jivraj S, Chee W, Corrado P (2006) Treatment planning of the edentulous maxilla. *Br Dent J* 201: 261-279.
- Choi TR, Jin TH, Dong JK (1995) A study on the exposure of maxillary and mandibular central incisor in smiling and physiologic rest position. *J Wonkwang Dent Res Inst* 5: 371-379.
- Neves FD, Mendonça G, Frenandes Neto AJ (2004) Analysis of influence of lip line and lip support in esthetics and selection of maxillary implant-supported prosthesis design. *J Prosthet Dent* 91(3): 286-288.
- Naert I, DeClercq M, Theuniers G, Schepers E (1988) Overdentures supported by osseointegrated fixtures for the edentulous mandible. A 2.5 year report. *Int J Oral Maxillofac Implants* 3(3): 191-196.
- Holden PK, Sufyan AS, Perkins SW (2011) Long-term analysis of surgical correction of the senile upper lip. *Arch Facial Plast Surg* 13(5): 332-336.
- Rifkin R (2000) Facial analysis: a comprehensive approach to treatment planning in aesthetic dentistry. *Pract Periodontics Aesthet Dent* 12(9): 865-871.
- Sabri R (2005) The eight components of a balanced smile. *J Clin Orthod* 39(3): 155-167.
- Wong WW, Davis DG, Camp MC, Gupta SC (2010) Contribution of lip proportions to facial aesthetics in different ethnicities: a three-dimensional analysis. *J Plast Reconstr Aesthet Surg* 63(12): 2032-2039.
- Niamtu J (2011) *Cosmetic Facial Surgery*. Elsevier, St. Louis, Missouri, USA, pp. 617-20, 709-716.
- Reyneke JP (2010) *Essentials of Orthognathic Surgery*. (2nd edn), Quintessence, Hanover Park, Illinois, USA, pp. 11-36.
- Zahrani AA (2007) Augmentation in two stages of atrophic alveolar bone prior to dental rehabilitation: a case report. *J Contemp Dent Pract* 8(6): 57-63.
- Kim SW, Park SO, Choi TH, Hai DT (2012) Change in upper lip height and nostril sill after alveolar bone grafting in unilateral cleft lip alveolus patients. *J Plast Reconstr Aesthet Surg* 65(5): 558-563.
- Mericske-Stern RD, Taylor TD, Belser U (2000) Management of the edentulous patient. *J Clin Oral Implants Res* 11(Suppl 1): 108-125.
- Salama M, Coachman C, Garber D, Calamita M, Salama H, et al. (2009) Prosthetic gingival reconstruction in the fixed partial restoration, part 2: diagnosis and treatment planning. *Int J Periodontics Restorative Dent* 29(6): 573-581.